

Homotopie et Homologie

Exercise Set 7

04.11.2010

Throughout these exercises, *space* means *topological space*, *map* means *continuous map*, and I denotes $[0, 1]$.

1. Let A be a subspace of X , and let $x_0 \in A$. Let P_i denote the homotopy fiber of the inclusion $i : A \hookrightarrow X$ (cf. Definition 2, Exercise Set 6). Show that

$$\pi_n(X, A) \cong \pi_{n-1}P_i.$$

2. Let $f : X \rightarrow Y$ be any pointed map, and let P_f denote the homotopy fiber of f . Prove that there is an exact sequence

$$\cdots \rightarrow \pi_n P_f \rightarrow \pi_n X \rightarrow \pi_n Y \rightarrow \pi_{n-1} P_f \rightarrow \cdots \rightarrow \pi_1 Y \rightarrow \pi_0 P_f \rightarrow \pi_0 X \rightarrow \pi_0 Y$$

and describe the homomorphisms in the sequence. This is the *long exact sequence in homotopy* of the *fiber sequence* $P_f \xrightarrow{qf} X \xrightarrow{f} Y$.

Hint 1. The existence of this long exact sequence can be proved using either the dual Barratt-Puppe sequence or the long exact sequence of a pair. Try to do it both ways!

3. Prove that if X is contractible, then $\pi_{n-1}P_f \cong \pi_n Y$ for all pointed maps $f : X \rightarrow Y$.
4. Let $\{x_0\} \subseteq A \subseteq B \subseteq X$ be a sequence of subspace inclusions. Show that there is an exact sequence

$$\cdots \rightarrow \pi_n(B, A) \rightarrow \pi_n(X, A) \rightarrow \pi_n(X, B) \xrightarrow{\partial_n} \pi_{n-1}(B, A) \rightarrow \cdots$$

Hint 2. The *connecting homomorphism* ∂_n is equal to the composite

$$\pi_n(X, B) \rightarrow \pi_{n-1}B \rightarrow \pi_{n-1}(B, A).$$

5. Consider the map $h : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ given by

$$h(x_1, x_2, x_3, x_4) = (x_1x_3 + x_2x_4, x_2x_3 - x_1x_4, x_1^2 + x_2^2 - x_3^2 - x_4^2).$$

- (a) Show that h restricts and corestricts to a map $h : S^3 \rightarrow S^2$. (This is the famous *Hopf map*.)
- (b) Show that the homotopy fiber P_h of h has the homotopy type of S^1 .
Hint 3. Start by showing that $h^{-1}(v) \cong S^1$ for any $v \in S^2$.
- (c) Show that $\pi_n S^1 = \{0\}$ for all $n > 1$.
Hint 4. Use the relation $\pi_n X \cong \pi_{n-1} \Omega X$.
- (d) It follows from the *Freudenthal suspension theorem* (to be proved later this semester) that

$$\pi_k S^n \cong \pi_{k+1} S^{n+1} \quad \forall k < 2n - 1.$$

Use this isomorphism to prove that $\pi_k S^n = \{0\}$ for all $1 \leq k < n$.

Hint 5. Recall that Seifert-van Kampen implies that S^n is simply connected for all $n \geq 2$.

- (e) Show that $\pi_2 S^2 \cong \mathbb{Z}$ (whence, by Freudenthal, $\pi_n S^n \cong \mathbb{Z}$ for all n) and that $\pi_3 S^2 \cong \mathbb{Z}$.
Hint 6. Use the long exact sequence in homotopy of h .